

"Express Mail" Mailing Label No. EL 847 787 600 US

Date of Deposit August 10, 2001

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Susan LaMont

Typed or Printed Name of Person Mailing Paper or Fee

Susan LaMont

Signature of Person Mailing Paper or Fee

Application for U.S. Letters Patent Entitled

IN UTERO ORAL NUCLEIC ACID IMMUNIZATION

claiming priority to U.S. Provisional Patent Application  
Serial No. 60,224,737, filed August 10, 2000

by Inventors:

Volker Gerdts  
Lorne Babiuk  
Sylvia van Drunen Littel-van den Hurk  
Philip J. Griebel

ROBINS & PASTERNAK LLP  
90 Middlefield Road, Suite 200  
Menlo Park, CA 94025  
Telephone: 650-325-7812  
Facsimile: 650-325-7823

Attorney Docket No. 9000-0058

5

IN UTERO ORAL NUCLEIC ACID IMMUNIZATION

Cross-Reference to Related Application

10           This application is related to provisional patent application serial no. 60/224,737, filed August 10, 2000, from which priority is claimed under 35 USC §119(e)(1) and which is incorporated herein by reference in its entirety.

Technical Field

15           The present invention relates generally to methods of nucleic acid immunization. In particular, the present invention pertains to methods of achieving immune responses by the *in utero* delivery of nucleic acid molecules that encode one or more selected antigens.

Background

20           Infectious diseases are the primary cause of neonatal morbidity and mortality in humans. The World Health Organization has estimated that in 1995 approximately 8 million (6.4% of live-born) infants died within the first year of life from these diseases, including 5 million during the first week of life. Some of the important pathogens involved include Herpes simplex virus (HSV), Human Immunodeficiency Virus (HIV),  
25   Hepatitis B virus (HBV), Human Cytomegalovirus (HCMV), Group B *Streptococcus* (GSB), *Haemophilus* and *Chlamydia* (Wright, et al., *Vaccine* (1998) 16:1355-1359; Mulholland, K., *Vaccine* (1998) 16:1360-1362). Infection with these pathogens can occur *in utero*, following early rupture of the amniotic membranes or during birth. In  
30   addition, infection may be transmitted during labor by non-sterile techniques, by breast feeding or during the first days of life in a perinatal nursery. To reduce the risk of disease transmission, caesarian sections, prophylactic treatment with antibiotics or maternal

antiviral therapy during the last trimester are used where available, together with improved neonatal care. None of these approaches, however, completely eliminates the risk of neonatal infection.

Since the first reports in 1993 (Ulmer, et al., *Science* (1993) 259:1745-1749), numerous studies have demonstrated that vaccination with DNA represents a very useful tool to induce immunity in people and animals (Donnelly, et al., *Annu Rev Immunol.* (1997) 15:617-648; Babiuk, et al., *Adv. Vet. Med.* (1999) 41:163-179. In addition to the simplicity of production and delivery, DNA vaccines possess the advantages of attenuated live vaccines with respect to their immunogenicity, and a level of biological safety similar to inactivated vaccines. DNA vaccines represent, therefore, a significant advance in vaccinology. Most studies with DNA vaccines have been performed in mature animals, but within the last three years, several studies have reported successful immunization of newborns of a variety of species (Butts, et al., *Vaccine* (1998) 16:1444-1449). As a result, DNA vaccines for rabies, hepatitis B, lymphocytic choriomeningitis-, influenza-, measles-, sendai-, porcine-, bovine herpesvirus-1, and tetanus toxoid are in development (Butts, et al., *supra*; Le Potier, et al., *Vet. Microbiol* (1997) 55:75-80; Van Drunen Littel-van den Hurk, et al., *Viral Immunology* (1999) 12:67-77). These approaches are designed, however, to prevent infections during the first weeks of life. In contrast, fetal immunization might prevent infection *in utero*, during birth and in the immediate postnatal period and may, therefore, have a significant impact on neonatal survival and the quality of life of infants.

A major factor for preventing an initial infection of the infant is the induction of effective mucosal immunity. This is of particular importance since the majority of infectious agents enter the host via the mucosal surfaces (Staats, et al., In *Mucosal Vaccines*, 1996 (Kiyono, H., et al., eds.) pp. 17-39). Newborns face an especially high risk of vertical disease transmission during birth and by breast feeding. Recent studies have shown that DNA vaccination via the mucosal surfaces can induce both mucosal and systemic immunity (McCluskie and Davis, *Crit. Rev. Immunol.* (1999) 19:303-329).

Watts, et al., *Nature Med.* (1999) 5:427-430 immunized fetal baboons against hepatitis B three times during the last trimester with 5µg of the recombinant hepatitis B surface antigen protein. Antigen-specific serum antibodies were detectable within 10

days after the second immunization in 75% (5/8) of the immunized fetuses. However, induction of a cell-mediated immune response was not evaluated. Sekhon and Larson, (*Nature. Med.* (1995) 11:1201-1203) demonstrated that introduction of adenoviral vectors into the amniotic fluid resulted in transgene expression in lung tissue of fetal rats.

- 5           The development of effective fetal immunization protocols would provide a valuable approach to reducing the high risk of diseases in newborn children.

#### Summary of the Invention

- 10           The present invention is based on the discovery that a single immunization with a nucleic acid vaccine, delivered orally to a fetus through the amniotic fluid, induces high serum antibody titers and cell-mediated immune responses in immunized fetuses. Moreover, this route of delivery provides direct access to a number of potential mucosal sites in the upper and lower respiratory tract and induces local immunity in the oral cavity. *In utero* nucleic acid immunization, therefore, provides an efficient means of
- 15           limiting vertical transmission of infectious diseases before, during and after birth.

- Accordingly, in one embodiment, the present invention provides a method of delivering a recombinant vector encoding a selected antigen to a fetal vertebrate subject *in utero* to elicit an immune response. The method comprises administering the recombinant vector orally via the amniotic fluid to the fetus, under conditions that permit
- 20           the expression of the antigen, thereby eliciting an immunological response to the antigen.

            In preferred embodiments, the vertebrate subject is a mammal. Additionally, the recombinant vector may be a nonviral vector.

- In additional embodiments, the invention is directed to a method of eliciting an immune response in a vertebrate subject. The method comprises administering a first
- 25           vaccine composition comprising a nucleic acid molecule encoding a selected antigen to a fetal vertebrate subject *in utero*, mucosally via the amniotic fluid, under conditions that permit the expression of the antigen, thereby eliciting an immunological response to the antigen.

- In certain embodiments, the nucleic acid molecule is included in a recombinant
- 30           vector. Moreover, the administering may be by nonviral-mediated delivery, the vertebrate subject may be a mammal and the antigen may be a viral antigen.



In another embodiment, the invention is directed to a method of eliciting an immune response in a vertebrate subject, wherein the method comprises administering a first vaccine composition comprising a recombinant vector encoding a selected antigen to a fetal vertebrate subject *in utero*, orally, via the amniotic fluid, under conditions that permit the expression of the antigen, thereby eliciting an immunological response to the antigen, wherein the administering is nonviral-mediated delivery.

In certain embodiments of the above-described methods, the viral antigen is a herpesvirus antigen. Moreover, delivery is done in the third trimester.

In additional embodiments, the invention further comprises administering a second vaccine composition to the vertebrate subject at birth to boost the immune response to the antigen encoded by the nucleic acid molecule in the first vaccine composition. The second vaccine composition may include the nucleic acid molecule present in the first vaccine composition. Alternatively, the second vaccine composition may be a subunit vaccine composition that includes the antigen encoded by the nucleic acid molecule present in the first vaccine composition.

In yet a further embodiment, the invention is directed to a method of eliciting an immune response in a mammalian subject. The method comprises:

(a) administering a first vaccine composition comprising a recombinant vector encoding a selected viral antigen to a fetal mammalian subject *in utero* during the third trimester, orally via the amniotic fluid, under conditions that permit the expression of the antigen and elicit an immunological response to the antigen; and

(b) administering a second vaccine composition to the mammalian subject at birth to boost the immune response to the antigen encoded by the recombinant vector in the first vaccine composition.

In certain embodiments, the second vaccine composition comprises the recombinant vector present in the first vaccine composition. In other embodiments, the second vaccine composition is a subunit vaccine composition that comprises the antigen encoded by the recombinant vector present in the first vaccine composition.

In yet other embodiments, the administering in step (a) and/or step (b) is nonviral-mediated delivery.

These and other embodiments of the subject invention will readily occur to those of skill in the art in view of the disclosure herein.

#### Brief Description of the Figures

- 5           Figure 1 shows lymphoproliferative responses of fetal blood mononuclear cells (PBMC) and retropharyngeal lymph node cells. Figure 1A shows the results of an experiment where PBMC from PBS (solid bar), tgD plasmid (grey bar), or irradiated BHV-1 (open bar) treated fetuses were stimulated with either 0.1  $\mu\text{g/ml}$  or 1.0  $\mu\text{g/ml}$  of recombinant BHV-1-tgD. Figure 1B shows the results of an experiment where
- 10   lymphocytes were isolated from fetal retropharyngeal lymph nodes of the same fetuses and stimulated with 0.1  $\mu\text{g/ml}$  or 1.0  $\mu\text{g/ml}$  of tgD-protein. The values presented are the mean of triplicate wells and are expressed as mean stimulation index (SI) for each group of treated fetuses. Error bars are one standard deviation. A statistically significant difference ( $p < 0.005$ ,  $t$ -test) between treatment groups is indicated (\*).
- 15           Figure 2 shows detection of gD-specific antibody secreting cells (ASC) in the retropharyngeal and right prescapular lymph nodes. The number of gD-specific ASC was quantified by incubating  $1 \times 10^6$  lymph node cells/well in triplicate wells that were either coated with tgD or PBS (negative control). Antigen-specific ASC were calculated by subtracting spots in control wells from spots in tgD-coated wells. Values represent the
- 20   average of two (2/4) DNA vaccinated fetuses (#5 and #6, grey bar), the BHV-1- (open bar) and the PBS-treated fetuses (black bar).
- Figure 3 shows serum IgG titers in blood samples from animals which had previously been immunized *in utero* with tgD (grey diamonds and black squares) or PBS (black triangles and black rectangles) and subsequently boosted at birth with tgD (grey
- 25   diamonds and black rectangles), or not boosted (black squares and black diamonds). The arrows represent the times immunizations were administered.
- Figure 4 shows lymphoproliferative responses of newborn blood mononuclear cells (PBMC) from tgD plasmid-boosted animals, as well as from animals which were not boosted.

Detailed Description

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Vols. I, II and III, Second Edition (1989); Perbal, B., *A Practical Guide to Molecular Cloning* (1984); the series, *Methods In Enzymology* (S. Colowick and N. Kaplan eds., Academic Press, Inc.); and *Handbook of Experimental Immunology*, Vols. I-IV (D.M. Weir and C.C. Blackwell eds., 1986, Blackwell Scientific Publications).

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

The following amino acid abbreviations are used throughout the text:

Alanine: Ala (A)	Arginine: Arg (R)
Asparagine: Asn (N)	Aspartic acid: Asp (D)
Cysteine: Cys (C)	Glutamine: Gln (Q)
Glutamic acid: Glu (E)	Glycine: Gly (G)
Histidine: His (H)	Isoleucine: Ile (I)
Leucine: Leu (L)	Lysine: Lys (K)
Methionine: Met (M)	Phenylalanine: Phe (F)
Proline: Pro (P)	Serine: Ser (S)
Threonine: Thr (T)	Tryptophan: Trp (W)
Tyrosine: Tyr (Y)	Valine: Val (V)

A. Definitions

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

It must be noted that, as used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “an antigen” includes a mixture of two or more antigens, and the like.

By "isolated" when referring to a nucleotide sequence, is meant that the indicated molecule is present in the substantial absence of other biological macromolecules of the same type. Thus, an "isolated" nucleic acid molecule which encodes a particular polypeptide refers to a nucleic acid molecule which is substantially free of other nucleic acid molecules that do not encode the subject polypeptide; however, the molecule may include some additional bases or moieties which do not deleteriously affect the basic characteristics of the composition.

For the purpose of describing the relative position of nucleotide sequences in a particular nucleic acid molecule throughout the instant application, such as when a particular nucleotide sequence is described as being situated "upstream," "downstream," "3'," or "5'" relative to another sequence, it is to be understood that it is the position of the sequences in the "sense" or "coding" strand of a DNA molecule that is being referred to as is conventional in the art.

The terms "polypeptide" and "protein" are used interchangeably herein and refer to a polymer of amino acid residues and are not limited to a minimum length of the product. Thus, peptides, oligopeptides, dimers, multimers, and the like, are included within the definition. Both full-length proteins and fragments thereof are encompassed by the definition. The terms also include postexpression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation and the like. Furthermore, for purposes of the present invention, a "polypeptide" refers to a protein which includes modifications, such as deletions, additions and substitutions (generally conservative in nature), to the native sequence, so long as the protein maintains the desired activity. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the proteins or errors due to PCR amplification.

The terms "analog" and "mutein" refer to biologically active derivatives of the reference molecule, or fragments of such derivatives, that retain desired activity, such as immunological activity as described herein. In general, the term "analog" refers to compounds having a native polypeptide sequence and structure with one or more amino acid additions, substitutions (generally conservative in nature) and/or deletions, relative to the native molecule, so long as the modifications do not destroy immunogenic activity.



The term "mutein" refers to peptides having one or more peptide mimics ("peptoids"), such as those described in International Publ. No. WO 91/04282. Preferably, the analog or mutein has at least the same immunoactivity as the native molecule. Methods for making polypeptide analogs and muteins are known in the art and are described further below.

Particularly preferred analogs include substitutions that are conservative in nature, i.e., those substitutions that take place within a family of amino acids that are related in their side chains. Specifically, amino acids are generally divided into four families: (1) acidic -- aspartate and glutamate; (2) basic -- lysine, arginine, histidine; (3) non-polar -- alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar -- glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. Phenylalanine, tryptophan, and tyrosine are sometimes classified as aromatic amino acids. For example, it is reasonably predictable that an isolated replacement of leucine with isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar conservative replacement of an amino acid with a structurally related amino acid, will not have a major effect on the biological activity. For example, the polypeptide of interest may include up to about 5-10 conservative or non-conservative amino acid substitutions, or even up to about 15-25 conservative or non-conservative amino acid substitutions, or any integer between 5-25, so long as the desired function of the molecule remains intact. One of skill in the art may readily determine regions of the molecule of interest that can tolerate change by reference to Hopp/Woods and Kyte-Doolittle plots, well known in the art.

By "antigen" is meant a molecule which contains one or more epitopes that will stimulate a host's immune system to make a cellular antigen-specific immune response when the antigen is produced, or a humoral antibody response. For purposes of the present invention, antigens can be derived from any of several known viruses, bacteria, parasites and fungi. Furthermore, for purposes of the present invention, an "antigen" refers to a protein which includes modifications, such as deletions, additions and substitutions (generally conservative in nature), to the native sequence, so long as the protein maintains the ability to elicit an immunological response. These modifications



may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the antigens.

By "fragment" is intended a polypeptide or polynucleotide consisting of only a part of the intact full-length polypeptide sequence and structure. An "immunogenic  
5 fragment" of a particular protein or will generally include at least about 5-10 contiguous amino acid residues of the full-length molecule, preferably at least about 15-25 contiguous amino acid residues of the full-length molecule, and most preferably at least about 20-50 or more contiguous amino acid residues of the full-length molecule, that define an epitope, or any integer between 5 amino acids and the full-length sequence,  
10 provided that the fragment in question retains the ability to elicit an immune response as defined below. A polynucleotide encoding such a fragment will include the requisite number of bases.

The term "epitope" as used herein refers to a sequence of at least about 3 to 5, preferably about 5 to 10 or 15, and not more than about 1,000 amino acids (or any integer  
15 therebetween), which define a sequence that by itself or as part of a larger sequence, will stimulate a host's immune system to make a cellular antigen-specific immune response when the antigen is presented, or a humoral antibody response. An epitope for use in the subject invention is not limited to a polypeptide having the exact sequence of the portion of the parent protein from which it is derived. Indeed, viral genomes are in a state of  
20 constant flux and contain several variable domains which exhibit relatively high degrees of variability between isolates. Thus the term "epitope" encompasses sequences identical to the native sequence, as well as modifications to the native sequence, such as deletions, additions and substitutions (generally conservative in nature).

Regions of a given polypeptide that include an epitope can be identified using any  
25 number of epitope mapping techniques, well known in the art. See, e.g., *Epitope Mapping Protocols* in *Methods in Molecular Biology*, Vol. 66 (Glenn E. Morris, Ed., 1996) Humana Press, Totowa, New Jersey. For example, linear epitopes may be determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the  
30 peptides with antibodies while the peptides are still attached to the supports. Such techniques are known in the art and described in, e.g., U.S. Patent No. 4,708,871; Geysen

et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:3998-4002; Geysen et al. (1986) *Molec. Immunol.* 23:709-715, all incorporated herein by reference in their entireties. Similarly, conformational epitopes are readily identified by determining spatial conformation of amino acids such as by, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., *Epitope Mapping Protocols, supra*. Antigenic regions of proteins can also be identified using standard antigenicity and hydropathy plots, such as those calculated using, e.g., the Omega version 1.0 software program available from the Oxford Molecular Group. This computer program employs the Hopp/Woods method, Hopp et al., *Proc. Natl. Acad. Sci USA* (1981) 78:3824-3828 for determining antigenicity profiles, and the Kyte-Doolittle technique, Kyte et al., *J. Mol. Biol.* (1982) 157:105-132 for hydropathy plots.

As used herein the term "T-cell epitope" refers to a feature of a peptide structure which is capable of inducing T-cell immunity towards the peptide structure or an associated hapten. T-cell epitopes generally comprise linear peptide determinants that assume extended conformations within the peptide-binding cleft of MHC molecules, (Unanue et al., *Science* (1987) 236:551-557). Conversion of polypeptides to MHC class II-associated linear peptide determinants (generally between 5 - 14 amino acids in length) is termed "antigen processing" which is carried out by antigen presenting cells (APCs). More particularly, a T-cell epitope is defined by local features of a short peptide structure, such as primary amino acid sequence properties involving charge and hydrophobicity, and certain types of secondary structure, such as helicity, that do not depend on the folding of the entire polypeptide. Further, it is believed that short peptides capable of recognition by helper T-cells are generally amphipathic structures comprising a hydrophobic side (for interaction with the MHC molecule) and a hydrophilic side (for interacting with the T-cell receptor), (Margalit et al., *Computer Prediction of T-cell Epitopes*, New Generation Vaccines Marcel-Dekker, Inc, ed. G.C. Woodrow et al., (1990) pp. 109-116) and further that the amphipathic structures have an  $\alpha$ -helical configuration (see, e.g., Spouge et al. *J. Immunol.* (1987) 138:204-212; Berkower et al. *J. Immunol.* (1986) 136:2498-2503).

Hence, segments of proteins which include T-cell epitopes can be readily predicted using numerous computer programs. (See e.g., Margalit et al., *Computer*

*Prediction of T-cell Epitopes*, New Generation Vaccines Marcel-Dekker, Inc, ed. G.C. Woodrow et al., (1990) pp. 109-116). Such programs generally compare the amino acid sequence of a peptide to sequences known to induce a T-cell response, and search for patterns of amino acids which are believed to be required for a T-cell epitope.

5           An "immunological response" to a polypeptide or composition is the development in a subject of a humoral and/or a cellular immune response to molecules present in the composition of interest. For purposes of the present invention, a "humoral immune response" refers to an immune response mediated by antibody molecules, while a "cellular immune response" is one mediated by T-lymphocytes and/or other white blood  
10       cells. One important aspect of cellular immunity involves an antigen-specific response by cytolytic T-cells ("CTLs"). CTLs have specificity for peptide antigens that are presented in association with proteins encoded by the major histocompatibility complex (MHC) and expressed on the surfaces of cells. CTLs help induce and promote the intracellular destruction of intracellular microbes, or the lysis of cells infected with such  
15       microbes. Another aspect of cellular immunity involves an antigen-specific response by helper T-cells. Helper T-cells act to help stimulate the function, and focus the activity of, nonspecific effector cells against cells displaying peptide antigens in association with MHC molecules on their surface. A "cellular immune response" also refers to the production of cytokines, chemokines and other such molecules produced by activated T-  
20       cells and/or other white blood cells, including those derived from CD4+ and CD8+ T-cells.

          A composition, such as an immunogenic composition, or vaccine that elicits a cellular immune response may serve to sensitize a vertebrate subject by the presentation of antigen in association with MHC molecules at the cell surface. The cell-mediated  
25       immune response is directed at, or near, cells presenting antigen at their surface. In addition, antigen-specific T-lymphocytes can be generated to allow for the future protection of an immunized host.

          The ability of a particular antigen or composition to stimulate a cell-mediated immunological response may be determined by a number of assays, such as by  
30       lymphoproliferation (lymphocyte activation) assays, CTL cytotoxic cell assays, or by assaying for T-lymphocytes specific for the antigen in a sensitized subject. Such assays

are well known in the art. See, e.g., Erickson et al., *J. Immunol.* (1993) 151:4189-4199; Doe et al., *Eur. J. Immunol.* (1994) 24:2369-2376; and the examples below.

Thus, an immunological response as used herein may be one which stimulates the production of CTLs, and/or the production or activation of helper T-cells. The antigen of interest may also elicit an antibody-mediated immune response. Hence, an immunological response may include one or more of the following effects: the production of antibodies by B-cells; and/or the activation of suppressor T-cells and/or  $\gamma\delta$  T-cells directed specifically to an antigen or antigens present in the composition or vaccine of interest. These responses may serve to neutralize infectivity, and/or mediate antibody-complement, or antibody dependent cell cytotoxicity (ADCC) to provide protection to an immunized host. Such responses can be determined using standard immunoassays and neutralization assays, well known in the art.

An "immunogenic" polypeptide or composition is one which elicits an immunological response as defined above.

The term "antibody" as used herein includes antibodies obtained from both polyclonal and monoclonal preparations, as well as, the following: hybrid (chimeric) antibody molecules (see, for example, Winter et al. (1991) *Nature* 349:293-299; and U.S. Patent No. 4,816,567); F(ab')<sub>2</sub> and F(ab) fragments; Fv molecules (non-covalent heterodimers, see, for example, Inbar et al. (1972) *Proc Natl Acad Sci USA* 69:2659-2662; and Ehrlich et al. (1980) *Biochem* 19:4091-4096); single-chain Fv molecules (sFv) (see, for example, Huston et al. (1988) *Proc Natl Acad Sci USA* 85:5879-5883); dimeric and trimeric antibody fragment constructs; minibodies (see, e.g., Pack et al. (1992) *Biochem* 31:1579-1584; Cumber et al. (1992) *J Immunology* 149B:120-126); humanized antibody molecules (see, for example, Riechmann et al. (1988) *Nature* 332:323-327; Verhoeyan et al. (1988) *Science* 239:1534-1536; and U.K. Patent Publication No. GB 2,276,169, published 21 September 1994); and, any functional fragments obtained from such molecules, wherein such fragments retain immunological binding properties of the parent antibody molecule.

A "recombinant" protein is a protein which retains the desired activity and which has been prepared by recombinant DNA techniques as described herein. In general, the gene of interest is cloned and then expressed in transformed organisms, as described



further below. The host organism expresses the foreign gene to produce the protein under expression conditions.

A "coding sequence" or a sequence which "encodes" a particular protein, is a nucleic acid sequence which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, viral nucleic acid sequences, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the coding sequence.

A "nucleic acid" sequence refers to a DNA or RNA sequence. The term captures sequences that include any of the known base analogues of DNA and RNA such as, but not limited to 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, -uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

The term DNA "control sequences" refers collectively to promoter sequences, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, and the like, which collectively provide for the replication, transcription and translation of a coding sequence in a recipient cell. Not all of these control sequences need always be

present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate host cell.

The term "promoter" is used herein in its ordinary sense to refer to a nucleotide region comprising a DNA regulatory sequence, wherein the regulatory sequence is derived from a gene which is capable of binding RNA polymerase and initiating transcription of a downstream (3'-direction) coding sequence. Transcription promoters can include "inducible promoters" (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), "repressible promoters" (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and "constitutive promoters".

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

A "vector" is a replicon, such as a plasmid, phage, or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. A vector is capable of transferring gene sequences to target cells (e.g., bacterial plasmid vectors, particulate carriers, and liposomes).

Typically, the terms "vector construct," "expression vector," "gene expression vector," "gene delivery vector," "gene transfer vector," and "expression cassette" all refer to an assembly which is capable of directing the expression of a sequence or gene of interest. Thus, the terms include cloning and expression vehicles.

By "nonviral-mediated delivery" is meant the use of a vector construct, as defined above, which does not use an infectious viral vector system to gain entry into the host cell. Thus, gene delivery techniques using retrovirus, adenovirus, sindbus virus, adeno-associated virus, and the like, are considered viral-mediated delivery techniques herein.

Generally, nonviral-mediated delivery refers to the use of a vector construct which includes the gene of interest operably linked to control elements which direct the expression of the gene of interest *in vivo*. The control elements can themselves be derived from viruses. One such nonviral-mediated delivery system involves the use of  
5 “naked” (plasmid) DNA. See, e.g., U.S. Patent Nos. 6,214,804; 5,910,488; 5,589,466; 5,580,859, all of which are incorporated herein by reference in their entireties.

A “host cell” is a cell which has been transformed, or is capable of transformation, by an exogenous nucleic acid molecule.

A cell has been “transformed” by exogenous DNA when such exogenous DNA  
10 has been introduced inside the cell membrane. Exogenous DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. In procaryotes and yeasts, for example, the exogenous DNA may be maintained on an episomal element, such as a plasmid. With respect to eucaryotic cells, a stably transformed cell is one in which the exogenous DNA has become integrated into the  
15 chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eucaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the exogenous DNA.

“Homology” refers to the percent similarity between two polynucleotide or two polypeptide moieties. Two DNA, or two polypeptide sequences are “substantially  
20 homologous” to each other when the sequences exhibit at least about 50% , preferably at least about 75%, more preferably at least about 80%-85%, preferably at least about 90%, and most preferably at least about 95%-98% sequence similarity over a defined length of the molecules. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence.

25 In general, “identity” refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Percent identity and similarity can be determined by a direct comparison of the sequence information between two molecules by aligning the sequences, counting the exact number of matches between the two aligned sequences, dividing by the length of  
30 the shorter sequence, and multiplying the result by 100. Readily available computer programs can be used to aid in the analysis, such as ALIGN, Dayhoff, M.O. in *Atlas of*



*Protein Sequence and Structure* M.O. Dayhoff ed., 5 Suppl. 3:353-358, National  
biomedical Research Foundation, Washington, DC, which adapts the local homology  
algorithm of Smith and Waterman *Advances in Appl. Math.* 2:482-489, 1981 for peptide  
analysis. Programs for determining nucleotide sequence similarity and identity are  
5 available in the Wisconsin Sequence Analysis Package, Version 8 (available from  
Genetics Computer Group, Madison, WI) for example, the BESTFIT, FASTA and GAP  
programs, which also rely on the Smith and Waterman algorithm. These programs are  
readily utilized with the default parameters recommended by the manufacturer and  
described in the Wisconsin Sequence Analysis Package referred to above. For example,  
10 percent identity of a particular nucleotide sequence to a reference sequence can be  
determined using the homology algorithm of Smith and Waterman with a default scoring  
table and a gap penalty of six nucleotide positions.

Another method of establishing percent identity or similarity in the context of the  
present invention is to use the MPSRCH package of programs copyrighted by the  
15 University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and  
distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the  
Smith-Waterman algorithm can be employed where default parameters are used for the  
scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a  
gap of six). From the data generated the "Match" value reflects "sequence identity."  
20 Other suitable programs for calculating the percent identity or similarity between  
sequences are generally known in the art, for example, another alignment program is  
BLAST, used with default parameters. For example, BLASTN and BLASTP can be used  
using the following default parameters: genetic code = standard; filter = none; strand =  
both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort  
25 by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB +  
GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs  
can be found at the following internet address: <http://www.ncbi.nlm.gov/cgi-bin/BLAST>.

Alternatively, homology can be determined by hybridization of polynucleotides  
under conditions which form stable duplexes between homologous regions, followed by  
30 digestion with single-stranded-specific nuclease(s), and size determination of the digested  
fragments. DNA sequences that are substantially homologous can be identified in a

Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

5 By "pharmaceutically acceptable" or "pharmacologically acceptable" is meant a material which is not biologically or otherwise undesirable, i.e., the material may be administered to an individual in a formulation or composition without causing any undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

10 By "nucleic acid immunization" is meant the introduction of a nucleic acid molecule encoding one or more selected antigens into a host, for the *in vivo* expression of an antigen, antigens, an epitope, or epitopes. The nucleic acid molecule can be introduced directly into a recipient subject, such as by mucosal delivery such as oral and intranasal administration, or the like, or can be introduced *ex vivo*, into cells which have  
15 been removed from the host. In the latter case, the transformed cells are reintroduced into the subject where an immune response can be mounted against the antigen encoded by the nucleic acid molecule.

By "mucosal" delivery is meant delivery of an antigen to a mucosal surface, including oral, nasal, pulmonary, vaginal, rectal, urethral, and sublingual or buccal  
20 delivery.

The term "treatment" as used herein refers to either (1) the prevention of infection or reinfection (prophylaxis), or (2) the reduction or elimination of symptoms of the disease of interest (therapy).

By the terms "vertebrate," "subject," and "vertebrate subject" are meant any  
25 member of the subphylum Chordata, including, without limitation, mammals such as cattle, sheep, pigs, goats, horses, and humans; domestic animals such as dogs and cats; and birds, including domestic, wild and game birds such as cocks and hens including chickens, turkeys and other gallinaceous birds; and fish. The term does not denote a particular age. Thus, both adult and newborn animals, as well as fetuses, are intended to  
30 be covered.

## B. Modes of Carrying Out the Invention

Before describing the present invention in detail, it is to be understood that this invention is not limited to particular formulations or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

Central to the present invention is the discovery that oral fetal immunization with a DNA vaccine elicits humoral and cellular immune responses, thus reducing the high risk of neonatal infections. Experiments performed in support of the present invention have clearly demonstrated that a single DNA immunization induced both humoral and cellular immune responses in all vaccinated animals. The observation that a single DNA injection induced active immunity has profound implications. In particular, the ability to avoid multiple immunizations clearly minimizes the risks associated with intrauterine injections. Moreover, the present results indicate that the fetal oral mucosa provides a very efficient site for DNA transfection.

The magnitude of the fetal immune responses in the present investigation was similar to that reported for 3 month old lambs following two intradermal injections of 500µg of the same plasmid (Van Drunen Littel-van den Hurk, et al., *Viral Immunology* (1999) 12:67-77). This is surprising, since it is well known that intradermal application of plasmid DNA induces significantly better systemic immune responses than mucosal or oral application of plasmid DNA (McCluskie and Davis, *Crit. Rev. Immunol.* (1999) 19:303-329; Donnelly, et al., *Annu Rev Immunol.* (1997) 15:617-648). Furthermore, delivery via the amniotic fluid provides a safe and relatively efficient method of DNA application that is effective without requiring formulation in exogenous transfection compounds.

The mucosal surfaces of the neonate are the primary site of entry for infectious agents during birth and breast feeding. Thus, the induction of both systemic and mucosal immunity provides optimal protection against most infectious diseases of newborns.



Recent observations suggest that both local and systemic immunity may contribute to immunity at mucosal surfaces (Baba, et al., *Nature Med.* (2000) 6:200-206; Mascola, et al., *Nature Med.* (2000) 6:207-210). In the examples detailed herein, mucosal immunity was evident in DNA-vaccinated fetuses through assays that detected gD-specific humoral (ASC) and cellular (LPR) immune responses in the lymph nodes draining the oral cavity. The presence of gD-specific serum antibody in all DNA-vaccinated fetuses was a clear indication of systemic immunity. Thus, *in utero*-DNA immunization during the last trimester induced both mucosal and systemic immune responses.

Mucosal immunity provides an important defense mechanism against a wide variety of pathogens. The mucosal surfaces of the gastrointestinal, respiratory and genitourinary tracts are continuously exposed to foreign antigens, including potentially infectious bacterial, viral and sometimes parasitic organisms. Mucosal immune responses protect against such challenges and have distinct and specialized characteristics.

For example, the principal immunoglobulin produced by the mucosal immune system is secretory IgA. Specialized antigen uptake cells in the Peyer's Patches of intestinal tract or nasopharyngeal lymphoid tissues, termed microfold or M cells, transport antigen to the underlying mucosal associated lymphoid tissues (MALT). In other areas of the mucosal epithelium, such as the pseudo-stratified airway epithelium, dendritic cells serve as antigen-presenting cells and migrate to local lymph nodes or MALT. Antigen processing and presentation occurs in the MALT, resulting in activation of antigen-specific IgA B cells. The subsequent trafficking and recirculation of the activated IgA B cells to other components of the mucosal immune system, e.g., the respiratory, intestinal and genital tracts, provides for disseminated local mucosal IgA responses throughout the "Common Mucosal System." Thus, the mucosal immune system is uniquely suited to respond to the types of antigenic challenge encountered by mucosal surfaces, and may provide the most effective type of immune response against particular pathogens. Accordingly, antigen delivery mechanisms which target the mucosal immune system provide an attractive means for achieving immunity.

A potential risk of immunizing fetuses and newborns is the induction of neonatal tolerance rather than the induction of an active immune response. However, since the

first reports of neonatal immunization in 1997 (Sarzotti, et al., *Vaccine* (1997) 15:795-797; Prince, et al., *Vaccine* (1997) 15:916-919), several studies have demonstrated successful DNA immunization in newborn animals (Butts, et al., *Vaccine* (1998) 16:1444-1449). Although the underlying mechanisms of the induction of neonatal

5 tolerance are still not completely understood, several parameters such as age, antigen dose and appropriate antigen presentation by dendritic cells might play an important role in developing neonatal nonresponsiveness (Ridge, et al., *Science* (1996) 271:1723-1726; Sarzotti, et al., *Science* (1996) 271:1726-1728; Forsthuber, et al., *Science* (1996) 271:1728-1730). So far, only immunization with the gene encoding the *Plasmodium*

10 *yoelii* malaria circumsporozoite protein (CSP) has induced neonatal tolerance in newborn mice (Mor, et al., *J. Clin. Invest.* (1996) 98:2700-2705; Ichino, et al., *J. Immunol.* (1999) 162:3814-3818; Isshii, et al., *Vaccine* (2000) 18:703-710). Induction of neonatal tolerance, as determined by the lack of a humoral or cellular response to a second DNA-injection after 6 weeks, was observed only after DNA immunization and not after

15 immunization with a recombinant protein (Mor, et al., *supra*). The development of tolerance also seemed to be age- (< 7days) and dose-dependent (10-100µg plasmid per mouse), but was not MHC-restricted (Ichino, et al., *supra*). Interestingly, CD8<sup>+</sup> T cells from tolerant mice transferred the induced nonresponsiveness to naïve recipients, which might indicate the involvement of suppressor T cells (Ichino, et al., *supra*). The same

20 investigators showed, however, that co-administration of a plasmid encoding the granulocyte-macrophage colony stimulating factor (GM-CSF) prevented the development of neonatal tolerance and elicited a primary IgG anti CSP immune response (Isshii, et al., *supra*) .

In contrast to this, the results provided herein clearly demonstrate that all fetuses

25 responded to the injection of plasmid-DNA and inactivated antigen with gD-specific IgG1-antibodies. The antibody titers induced in fetuses were higher than those observed following intradermal immunization of neonates with the same dose of the same plasmid. DNA immunized neonates also responded to a secondary DNA immunization at a later date (Van Drunen Littel-van den Hurk, et al., *Viral Immunology* (1999) 12:67-77). These

30 observations support the conclusion that DNA immunization of the late term fetus with

an antigen-encoding gene, such as the herpesvirus gD gene, should not induce neonatal tolerance.

DNA vaccination provides several inherent advantages (Donnelly, et al., *Annu Rev Immunol.* (1997) 15:617-648). The endogenous synthesis of an encoded antigen facilitates both MHC-I and MHC-II restricted antigen presentation, which is necessary for the *in vivo* generation of both cell-mediated and humoral immune responses. For example, in the examples detailed herein, DNA-vaccination was clearly more efficient than immunization with the corresponding purified antigen. All DNA-vaccinated fetuses displayed higher serum antibody titers and antigen-specific proliferative responses in blood and retropharyngeal lymph nodes, and there was no detectable mucosal immunity following immunization with inactivated virus (Table 1, Figures 1A and 1B). The endogenous synthesis of encoded proteins also ensures appropriate processing and conformation of viral proteins, which is particularly important for the generation of neutralizing antibodies (Van Drunen Littel-van den Hurk, et al., *Vaccine* (1992) 11:25-35). Previous studies have demonstrated that for alphaherpesviruses the conformation of the glycoprotein gD is a critical parameter for the induction of protective immunity (Van Drunen Littel-van den Hurk, et al., *Vaccine* (1992) 11:25-35). The appropriate expression of glycoprotein D in immunized fetuses was confirmed herein by the induction of neutralizing serum antibodies (Table 1). Thus, DNA vaccination of the fetus induced a humoral immune response that should protect against viral infection.

Unlike viral vectors, DNA vaccines can be injected repeatedly to effectively boost the level of an immune response (Donnelly, et al., *Annu Rev Immunol.* (1997) 15:617-648). This is of particular importance, since Iwamoto et al., *Gene Ther.* (1999) 6:98-106, recently showed that the repeated *in utero* delivery of an E1,E3-deleted adenoviral vector, encoding  $\beta$ -galactosidase, resulted in over 50% mortality in fetal lambs. Therefore, a second vaccination at birth, even with other types of vaccines, such as a subunit or modified live vaccines, could enhance the level of disease protection, acquired by neonates, who face a high risk of disease transmission. Moreover, a primary fetal immunization combined with secondary immunization at birth may confer protective immunity against the typical "first week-infections", such as human respiratory

syncytical virus (Henderson, F.W. in *Rudolph's Pediatrics* 1996 (Rudolph, A.M., et al., eds.) pp. 536-544, Appleton & Lange, Stamford, Connecticut).

Fetal sheep, the animal model used herein, provide an excellent model for studying several aspects of the ontogeny of immunity (Hein, W.R., *The Immunologist* (1995) 3:12-18) as well as new approaches for gene therapy (Zanjani, et al., *Science* (1999) 285:2084-2088) and human stem cells transplantation (Sekhon and Larson, *Nature. Med.* (1995) 11:1201-1203). In contrast to mice and rats, where the immune system undergoes very little peripheral expansion prior to birth, the peripheral lymphoid organs of sheep and humans undergo substantial development during fetal life (Griebel and Hein *Immunology Today* (1996) 17:30-39). Furthermore, sheep and humans show significant similarities in the appearance of leukocytes in circulation during fetal development (Alsalami, et al., *Aust. Vet. J.* (1999) 77:588-594). Immune responsiveness develops in fetal lambs by mid-gestation, and depending on the type of antigen, the capacity to perform isotype-switching has developed prior to the last trimester of gestation (Griebel, P.J. in *Handbook of Vertebrate Immunology* 1998 (Pastoret, et al., eds) pp. 485-554 (Academic Press, San Diego/London). In addition, the size of the fetal lamb during the last trimester of gestation facilitates a variety of experimental interventions. Thus, fetal sheep provide an appropriate model for investigating *in utero* immunization and making predictions from the results obtained using this animal model.

In order to further an understanding of the invention, a more detailed discussion is provided below regarding particular antigens for use with the subject methods, as well as methods of obtaining and delivering the compositions.

In particular, as explained above, the method of the invention results in both cell-mediated immunity as well as humoral antibody responses. Thus, in addition to a conventional antibody response, the system herein described provides for, e.g., the association of the expressed antigens with class I MHC molecules such that an *in vivo* cellular immune response to the antigen of interest can be mounted which stimulates the production of CTLs to allow for future recognition of the antigen. Furthermore, the method is capable of eliciting an antigen-specific response by helper T-cells.

Accordingly, the methods of the present invention will find use with any antigen for which both cellular and humoral immune responses are desired, including antigens

derived from viral, bacterial, fungal and parasitic pathogens that may induce antibodies, T-cell helper epitopes and T-cell cytotoxic epitopes. Such antigens include, but are not limited to, those encoded by human and animal viruses and can correspond to either structural or non-structural proteins.

5           The technique is particularly useful for immunization against intracellular viruses which normally elicit poor immune responses. Additionally, immunization against pathogens that threaten newborns is particularly desirable. Thus, the methods described herein will find use for immunizing against a wide variety of organisms, such as viral, bacterial, mycoplasma, parasitic and even tumor antigens, as well as allergens.

10           For example, the present invention will find use for stimulating an immune response against a wide variety of proteins from the herpesvirus family, including proteins derived from bovine herpes virus (BHV) and human herpes simplex virus (HSV) types 1 and 2, such as BHV-1, BHV-2, HSV-1 and HSV-2 glycoproteins gB, gD and gH; antigens derived from varicella zoster virus (VZV), Epstein-Barr virus (EBV) and  
15           cytomegalovirus (CMV) including CMV gB and gH; and antigens derived from other human and animal herpesviruses such as HHV6 and HHV7. (See, e.g. Chee et al., *Cytomegaloviruses* (J.K. McDougall, ed., Springer-Verlag 1990) pp. 125-169, for a review of the protein coding content of cytomegalovirus; McGeoch et al., *J. Gen. Virol.* (1988) 69:1531-1574, for a discussion of the various HSV-1 encoded proteins; U.S.  
20           Patent No. 5,171,568 for a discussion of HSV-1 and HSV-2 gB and gD proteins and the genes encoding therefor; Baer et al., *Nature* (1984) 310:207-211, for the identification of protein coding sequences in an EBV genome; and Davison and Scott, *J. Gen. Virol.* (1986) 67:1759-1816, for a review of VZV.) Additionally, immunization against various hepatitis viruses, such as HBV, using, for example, hepatitis B surface antigen (HBsAg),  
25           as well as any of the various human papilloma viruses, rotaviruses, etc., will find use with the present methods.

          Polynucleotide sequences encoding antigens derived from other viruses will also find use in the claimed methods, such as without limitation, proteins from members of the families Picornaviridae (e.g., polioviruses, etc.); Caliciviridae; Togaviridae (e.g., rubella  
30           virus, dengue virus, etc.); Flaviviridae; Coronaviridae; Reoviridae; Birnaviridae; Rhabdoviridae (e.g., rabies virus, etc.); Filoviridae; Paramyxoviridae (e.g., mumps



virus, measles virus, respiratory syncytial virus, etc.); Orthomyxoviridae (e.g., influenza virus types A, B and C, etc.); Bunyaviridae; Arenaviridae; Retroviridae (e.g., HTLV-I; HTLV-II; HIV-1 (also known as HTLV-III, LAV, ARV, hTLR, etc.)), including but not limited to antigens from the isolates HIV<sub>IIIb</sub>, HIV<sub>SF2</sub>, HIV<sub>LAV</sub>, HIV<sub>LAI</sub>, HIV<sub>MN</sub>; HIV-1<sub>CM235</sub>, HIV-1<sub>US4</sub>; HIV-2; simian immunodeficiency virus (SIV) among others. See, e.g. Virology, 3rd Edition (W.K. Joklik ed. 1988); *Fundamental Virology*, 2nd Edition (B.N. Fields and D.M. Knipe, eds. 1991), for a description of these and other viruses.

Antigens derived from fungal pathogens, bacterial pathogens, such as from *Mycobacterium leprae*, *M. tuberculosis*, *Borrelia burgdorferi*, *Bordetella pertussis*, *Clostridium tetani*, *Streptococcus*, *Salmonella typhimurium*, *S. enterica*, *Shigella*, *E. coli*, *Hemophilus influenzae*, and the like will also find use herein. Similarly, antigens from *Mycoplasma*, such as *M. pulmonis*; parasitic antigens, such as from *Plasmodium yoelii*, *P. falciparum*, *Toxoplasma gondii*, *Schistosoma japonicum*, *Leishmania major*, *Trypanosoma cruzi*, and so forth, can be used with the subject methods.

The techniques can be used for the delivery of discrete antigens, larger portions of the genome in question and, for example, a proviral DNA which includes nearly all of the viral genome. It is readily apparent that the subject invention can be used to prevent or treat a wide variety of diseases.

Additionally, rather than delivering the nucleic acid sequences encoding the antigens, nucleic acid sequences encoding recombinant antibodies may also be delivered by the subject methods.

Polynucleotide sequences coding for the above-described molecules can be obtained using recombinant methods, such as by screening cDNA and genomic libraries from cells expressing the gene, or by deriving the gene from a vector known to include the same. Furthermore, the desired gene can be isolated directly from cells and tissues containing the same, using standard techniques, such as phenol extraction and PCR of cDNA or genomic DNA. See, e.g., Sambrook et al., *supra*, for a description of techniques used to obtain and isolate DNA. The gene of interest can also be produced synthetically, rather than cloned. The nucleotide sequence can be designed with the appropriate codons for the particular amino acid sequence desired. In general, one will select preferred codons for the intended host in which the sequence will be expressed.

The complete sequence is assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence. See, e.g., Edge, *Nature* (1981) 292:756; Nambair et al., *Science* (1984) 223:1299; Jay et al., *J. Biol. Chem.* (1984) 259:6311.

5           Next, the gene sequence encoding the desired antigen can be inserted into a vector which includes control sequences operably linked to the desired coding sequence, which allow for the expression of the gene *in vivo* in the subject species. For example, typical promoters for mammalian cell expression include the SV40 early promoter, a CMV promoter such as the CMV immediate early promoter, the mouse mammary tumor virus  
10 LTR promoter, the adenovirus major late promoter (Ad MLP), and the herpes simplex virus promoter, among others. Other nonviral promoters, such as a promoter derived from the murine metallothionein gene, will also find use for mammalian expression. Typically, transcription termination and polyadenylation sequences will also be present, located 3' to the translation stop codon. Preferably, a sequence for optimization of  
15 initiation of translation, located 5' to the coding sequence, is also present. Examples of transcription terminator/polyadenylation signals include those derived from SV40, as described in Sambrook et al., *supra*, as well as a bovine growth hormone terminator sequence. Introns, containing splice donor and acceptor sites, may also be designed into the constructs for use with the present invention.

20           Enhancer elements may also be used herein to increase expression levels of the mammalian constructs. Examples include the SV40 early gene enhancer, as described in Dijkema et al., *EMBO J.* (1985) 4:761, the enhancer/promoter derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus, as described in Gorman et al., *Proc. Natl. Acad. Sci. USA* (1982b) 79:6777 and elements derived from human CMV, as  
25 described in Boshart et al., *Cell* (1985) 41:521, such as elements included in the CMV intron A sequence.

          Furthermore, plasmids can be constructed which include a chimeric gene sequence, encoding e.g., multiple antigens of interest, for example derived from more than one viral isolate. Additionally, genes coding for immune modulating agents which  
30 can enhance antigen presentation, attract lymphocytes to the site of gene expression or promote expansion of the population of lymphocytes to the site of gene expression or

promote expansion of the population of lymphocytes which respond to the expressed antigen, can also be present. Such agents include cytokines, lymphokines, and chemokines, including but not limited to IL-2, modified IL-2 (cys125-ser125), GM-CSF, IL-12,  $\gamma$ -interferon, IP-10, MIP1 $\alpha$ , MIP1 $\beta$  and RANTES. Additionally, immune molecules such as TAP transporters, costimulatory molecules such as B7,  $\beta$ 2M, class I or II MHC genes (syngeneic or allogeneic), and other genes coding for proteins that are required for efficient immune responses but are not expressed due to specific inhibition or deletion, will also find use in the constructs.

The above sequences can be administered using separate vectors or can be present on the vector bearing the gene encoding the antigen of interest. If present on the same vector, the additional gene sequences can either precede or follow the gene encoding the antigen of interest in a dicistronic gene configuration. Additional control elements can be situated between the various genes for efficient translation of RNA from the distal coding region. Alternatively, a chimeric transcription unit having a single open reading frame encoding both the gene of interest and the modulator, can also be constructed. Either a fusion can be made to allow for the synthesis of a chimeric protein or alternatively, protein processing signals can be engineered to provide cleavage by a protease such as a signal peptidase, thus allowing liberation of the two or more proteins derived from translation of the template RNA. Such signals for processing of a polyprotein exist in, e.g., flaviviruses, pestiviruses such as HCV, and picornaviruses, and can be engineered into the constructs. The processing protease may also be expressed in this system either independently or as part of a chimera with the antigen and/or cytokine coding region(s). The protease itself can be both a processing enzyme and a vaccine antigen.

Once complete, the constructs are used for nucleic acid immunization using standard gene delivery protocols. Methods for gene delivery are known in the art. See, e.g., U.S. Patent Nos. 5,399,346, 5,580,859, 5,589,466. Genes can be delivered either directly via the amniotic fluid to the vertebrate subject or, alternatively, delivered *ex vivo*, to cells derived from the subject and the transformed cells delivered to the subject. Moreover, the genes can be delivered with or without a viral vector.

A number of viral based systems have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene

delivery systems. A selected gene can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to cells of the subject either *in vivo* or *ex vivo*. A number of retroviral systems have been described (U.S. Patent No. 5,219,740; Miller and Rosman,

5 *BioTechniques* (1989) 7:980-990; Miller, A.D., *Human Gene Therapy* (1990) 1:5-14; Scarpa et al., *Virology* (1991) 180:849-852; Burns et al., *Proc. Natl. Acad. Sci. USA* (1993) 90:8033-8037; and Boris-Lawrie and Temin, *Cur. Opin. Genet. Develop.* (1993) 3:102-109.

A number of adenovirus vectors have also been described. Unlike retroviruses  
10 which integrate into the host genome, adenoviruses persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham, *J. Virol.* (1986) 57:267-274; Bett et al., *J. Virol.* (1993) 67:5911-5921; Mittereder et al., *Human Gene Therapy* (1994) 5:717-729; Seth et al., *J. Virol.* (1994) 68:933-940; Barr et al., *Gene Therapy* (1994) 1:51-58; Berkner, K.L. *BioTechniques* (1988) 6:616-629; and  
15 Rich et al., *Human Gene Therapy* (1993) 4:461-476).

Additionally, various adeno-associated virus (AAV) vector systems have been developed for gene delivery. AAV vectors can be readily constructed using techniques well known in the art. See, e.g., U.S. Patent Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 (published 23 January 1992) and WO 93/03769  
20 (published 4 March 1993); Lebkowski et al., *Molec. Cell. Biol.* (1988) 8:3988-3996; Vincent et al., *Vaccines 90* (1990) (Cold Spring Harbor Laboratory Press); Carter, B.J. *Current Opinion in Biotechnology* (1992) 3:533-539; Muzyczka, N. *Current Topics in Microbiol. and Immunol.* (1992) 158:97-129; Kotin, R.M. *Human Gene Therapy* (1994) 5:793-801; Shelling and Smith, *Gene Therapy* (1994) 1:165-169; and Zhou et al., *J. Exp.*  
25 *Med.* (1994) 179:1867-1875.

A vaccinia based infection/transfection system can be conveniently used to provide for inducible, transient expression of the gene of interest in a host cell. In this system, cells are first infected *in vitro* with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in  
30 that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the polynucleotide of interest, driven by a T7 promoter. The polymerase

expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into protein by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation products. See, e.g., Elroy-Stein and Moss, 5 *Proc. Natl. Acad. Sci. USA* (1990) 87:6743-6747; Fuerst et al., *Proc. Natl. Acad. Sci. USA* (1986) 83:8122-8126.

As an alternative approach to infection with vaccinia or avipox virus recombinants, or to the delivery of genes using other viral vectors, an amplification system can be used that will lead to high level expression following introduction into host 10 cells. Specifically, a T7 RNA polymerase promoter preceding the coding region for T7 RNA polymerase can be engineered. Translation of RNA derived from this template will generate T7 RNA polymerase which in turn will transcribe more template. Concomitantly, there will be a cDNA whose expression is under the control of the T7 promoter. Thus, some of the T7 RNA polymerase generated from translation of the 15 amplification template RNA will lead to transcription of the desired gene. Because some T7 RNA polymerase is required to initiate the amplification, T7 RNA polymerase can be introduced into cells along with the template(s) to prime the transcription reaction. The polymerase can be introduced as a protein or on a plasmid encoding the RNA polymerase. For a further discussion of T7 systems and their use for transforming cells, 20 see, e.g., International Publication No. WO 94/26911; Studier and Moffatt, *J. Mol. Biol.* (1986) 189:113-130; Deng and Wolff, *Gene* (1994) 143:245-249; Gao et al., *Biochem. Biophys. Res. Commun.* (1994) 200:1201-1206; Gao and Huang, *Nuc. Acids Res.* (1993) 21:2867-2872; Chen et al., *Nuc. Acids Res.* (1994) 22:2114-2120; and U.S. Patent No. 5,135,855.

25 The gene of interest can also be delivered without a viral vector. For example, the gene can be delivered directly to the subject or packaged in liposomes prior to delivery to the subject or to cells derived therefrom. Lipid encapsulation is generally accomplished using liposomes which are able to stably bind or entrap and retain nucleic acid. The ratio of condensed DNA to lipid preparation can vary but will generally be around 1:1 (mg 30 DNA:micromoles lipid), or more of lipid. For a review of the use of liposomes as carriers

for delivery of nucleic acids, see, Hug and Sleight, *Biochim. Biophys. Acta.* (1991) 1097:1-17; Straubinger et al., in *Methods of Enzymology* (1983), Vol. 101, pp. 512-527.

Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations, with cationic liposomes particularly preferred. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., *Proc. Natl. Acad. Sci. USA* (1987) 84:7413-7416); mRNA (Malone et al., *Proc. Natl. Acad. Sci. USA* (1989) 86:6077-6081); and purified transcription factors (Debs et al., *J. Biol. Chem.* (1990) 265:10189-10192), in functional form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, NY. (See, also, Felgner et al., *Proc. Natl. Acad. Sci. USA* (1987) 84:7413-7416). Other commercially available lipids include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer). Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g., Szoka et al., *Proc. Natl. Acad. Sci. USA* (1978) 75:4194-4198; PCT Publication No. WO 90/11092 for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, AL), or can be easily prepared using readily available materials. Such materials include phosphatidyl choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios.

Methods for making liposomes using these materials are well known in the art.

The liposomes can comprise multilammellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs). The various liposome-nucleic acid complexes are prepared using methods known in the art. See, e.g., Straubinger et al., in *METHODS OF IMMUNOLOGY* (1983), Vol. 101, pp. 512-527; Szoka et al., *Proc. Natl. Acad. Sci. USA* (1978) 75:4194-4198; Papahadjopoulos et al., *Biochim. Biophys. Acta* (1975) 394:483; Wilson et al., *Cell* (1979) 17:77; Deamer and Bangham, *Biochim.*

*Biophys. Acta* (1976) 443:629; Ostro et al., *Biochem. Biophys. Res. Commun.* (1977) 76:836; Fraley et al., *Proc. Natl. Acad. Sci. USA* (1979) 76:3348); Enoch and Strittmatter, *Proc. Natl. Acad. Sci. USA* (1979) 76:145); Fraley et al., *J. Biol. Chem.* (1980) 255:10431; Szoka and Papahadjopoulos, *Proc. Natl. Acad. Sci. USA* (1978) 75:145; and  
5 Schaefer-Ridder et al., *Science* (1982) 215:166.

The DNA can also be delivered in cochleate lipid compositions similar to those described by Papahadjopoulos et al., *Biochem. Biophys. Acta.* (1975) 394:483-491. See, also, U.S. Patent Nos. 4,663,161 and 4,871,488.

The gene of interest may also be delivered using particulate systems and  
10 polymers. For example, polymers such as polylysine, polyarginine, polyornithine, spermine, spermidine, as well as conjugates of these molecules, are useful for transferring a nucleic acid of interest. Similarly, DEAE dextran-mediated transfection, calcium phosphate precipitation or precipitation using other insoluble inorganic salts, such as strontium phosphate, aluminum silicates including bentonite and kaolin, chromic oxide,  
15 magnesium silicate, talc, and the like, will find use with the present methods. See, e.g., Felgner, P.L., *Advanced Drug Delivery Reviews* (1990) 5:163-187, for a review of delivery systems useful for gene transfer.

Additionally, biolistic delivery systems employing particulate carriers such as gold and tungsten, are especially useful for delivering genes of interest. The particles are  
20 coated with the gene to be delivered and accelerated to high velocity, generally under a reduced atmosphere, using a gun powder discharge from a "gene gun." For a description of such techniques, and apparatuses useful therefore, see, e.g., U.S. Patent Nos. 4,945,050; 5,036,006; 5,100,792; 5,179,022; 5,371,015; and 5,478,744.

The recombinant vectors (with or without associated lipids or carriers) are  
25 formulated into compositions for delivery to the vertebrate subject. Methods of preparing such formulations are described in, e.g., *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, Pennsylvania, 18 Edition, 1990. These compositions may either be prophylactic (to prevent infection) or therapeutic (to treat disease after infection). The compositions will comprise a "therapeutically effective amount" of the  
30 gene of interest such that an amount of the antigen can be produced *in vivo* so that an immune response is generated in the individual to which it is administered. The exact

amount necessary will vary depending on the subject being treated; the age and general condition of the subject to be treated; the capacity of the subject's immune system to synthesize antibodies; the degree of protection desired; the severity of the condition being treated; the particular antigen selected and its mode of administration, among other factors. An appropriate effective amount can be readily determined by one of skill in the art. Thus, a "therapeutically effective amount" will fall in a relatively broad range that can be determined through routine trials. For example, for purposes of the present invention, an effective dose will be from about 1  $\mu$ g to about 100 mg, more preferably from about 10  $\mu$ g to about 1 mg, of the DNA constructs.

The compositions will generally include one or more "pharmaceutically acceptable excipients or vehicles" appropriate for mucosal delivery, such as water, saline, glycerol, polyethyleneglycol, hyaluronic acid, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. Certain facilitators of nucleic acid uptake and/or expression can also be included in the compositions or coadministered, such as, but not limited to, bupivacaine, cardiotoxin and sucrose.

For example, intranasal and pulmonary formulations will usually include vehicles that neither cause irritation to the nasal mucosa nor significantly disturb ciliary function. Diluents such as water, aqueous saline or other known substances can be employed with the subject invention. The nasal formulations may also contain preservatives such as, but not limited to, chlorobutanol and benzalkonium chloride. A surfactant may be present to enhance absorption of the subject proteins by the nasal mucosa.

For rectal and urethral suppositories, the vehicle composition will include traditional binders and carriers, such as, cocoa butter (theobroma oil) or other triglycerides, vegetable oils modified by esterification, hydrogenation and/or fractionation, glycerinated gelatin, polyalkaline glycols, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

For vaginal delivery, the formulations of the present invention can be incorporated in pessary bases, such as those including mixtures of polyethylene triglycerides, or suspended in oils such as corn oil or sesame oil, optionally containing colloidal silica. See, e.g., Richardson et al., *Int. J. Pharm.* (1995) 115:9-15.



For a further discussion of appropriate vehicles to use for particular modes of delivery, see, e.g., *Remington: The Science and Practice of Pharmacy*, Mack Publishing Company, Easton, Pennsylvania, 19th edition, 1995. One of skill in the art can readily determine the proper vehicle to use for the particular antigen and site of delivery.

5           Once formulated, the compositions of the invention can be administered directly to the subject or, alternatively, delivered *ex vivo*, to cells derived from the subject, using methods such as those described above. For example, methods for the *ex vivo* delivery and reimplantation of transformed cells into a subject are known in the art and will include e.g., dextran-mediated transfection, calcium phosphate precipitation, polybrene  
10       mediated transfection, lipofectamine and LT-1 mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

          Direct delivery of the compositions *in vivo* will generally be accomplished with or without viral vectors, as described above, by injection using a conventional syringe or a  
15       gene gun, such as the Accell® gene delivery system (Agracetus, Inc., Middleton, WI). The constructs are injected mucosally, such as orally, via the amniotic fluid. The constructs may be delivered into cells of the epidermis which provides access to skin-associated lymphoid cells and provides for a transient presence of DNA in the vaccine recipient.

20           Dosage treatment may be a single dose schedule or a multiple dose schedule. For example, the first vaccine is normally given after the fetal immune system has developed to a point that renders it capable of developing an immune response and development has progressed to a stage that makes administration feasible. The organs which function in the immune system need not be completely developed at the time of the first vaccination.  
25       The time for delivery will depend on the particular species of subject in question. Thus, the DNA vaccines described herein will normally be delivered at least half-way through the gestational period, more preferably in the last trimester, but can be delivered earlier.

          As shown herein, immune responses are boostable with one or more injections given at birth or shortly thereafter, such as at birth or 1 day to 14 days or more after birth.  
30       Subsequent boosts can also be administered. The subsequent vaccine can be the same or different than the vaccine delivered *in utero*. Thus, the identical nucleic acid vaccine can



be delivered. Alternatively, a subunit vaccine containing the antigen encoded by the nucleic acid can be delivered, or even an attenuated bacterium or virus can be used for the boost.

5 C. Experimental

Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way. Those of skill in the art will readily appreciate that the invention may be practiced in a variety of ways given the  
10 teaching of this disclosure.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

15 Materials and Methods

In the following methods, restriction and modifying enzymes, as well as other reagents for DNA manipulations were purchased from commercial sources, and used according to the manufacturers' directions. In the cloning of DNA fragments, except  
20 where noted, all DNA manipulations were done according to standard procedures. See, e.g., Sambrook et al., *supra*.

**Plasmids:** Generation of the pSLIA<sub>gD</sub> plasmid was described previously by Braun *et al.* (*Vaccine Res.* (1997) 6:151-164). The gene encoding the truncated version  
25 of BHV-1 gD was inserted into plasmid pSLIA0 under the control of the human cytomegalovirus (HCMV) immediate early promoter/1A-region. Plasmids were transformed into *E. coli* DH5 $\alpha$  and purified using the Qiagen Giga-Kit (Qiagen, Mississauga, ON). DNA concentration was estimated after agarose gel electrophoresis by comparison with known standards and by spectrophotometry. For immunization, DNA  
30 was dissolved in pyrogen-free PBS, pH 7.3.

**Inactivation of BHV-1 by UV-irradiation:** BHV-1 was inactivated by UV-irradiation as described previously (Popowych, Y.I. An assessment of in vitro assays used to detect BHV-1 specific CMI responses. (1994) *Thesis, Department of Veterinary Microbiology, University of Saskatchewan, Saskatoon*). Briefly, one ml of  $1 \times 10^8$  p.f.u. of BHV-1 was inactivated in a cabinet fitted with a germicidal fluorescent bulb (G8T5, General Electronic) with a UV-dose of  $3260 \text{ mJ/cm}^2$ . Complete inactivation was verified in a virus-plaque assay.

**Fetal immunization:** Suffolk sheep were obtained from the Department of Animal and Poultry Science, University of Saskatchewan. The ewes were bred following estrus synchronization with medoxyprogesterone acetate (Veramix, Upjohn, Orangeville, ON) and injection with pregnant mare serum gonadotrophin (Equinex, Ayerst, Winnipeg, MB). Pregnancy was confirmed by ultrasound at day 45 and 105 of gestation and fetal immunization was then performed at days 123 or 124 of gestation. Briefly, ewes were anaesthetized as described previously (Mutwiri, et al., *Immunology* (1999) 97:455-461) and positioned in dorsal recumbancy with a slight rotation to the right side. A ventral midline incision was made in the abdomen and both uterine horns were palpated to verify the presence of a single or two fetuses. Subsequently, the head of the fetus was positioned with the uterine wall compressed against the commissure of the fetal mouth. The fetal mouth was opened and 5 ml of DNA-solution or sterile PBS were injected using a 23g-needle inserted through the uterine wall and positioned in the fetal oral cavity. This injection was followed with 500 mg of ampicillin (250mg/ml; Novopharm, Toronto, ON). For injection of the inactivated BHV-1, the fetus was positioned with its right side underlying the uterine wall. A 2 cm incision was made through the uterine wall, without disrupting the amniotic membrane. A purse-string suture with 4-0 PDS (Ethicon, Peterborough, ON) was placed in the amniotic membrane at the margins of the uterine incision. The amniotic membrane was incised in the centre of the purse-string suture and forceps were used to raise a fold of skin on the neck of the fetus. One ml each of irradiated BHV-1 ( $2.5 \times 10^7$  p.f.u.) was injected subcutaneously at the base of the skinfold and into the oral cavity followed by a second oral injection of 500mg ampicillin. The purse-string suture was used to close the amniotic membranes and the uterine wall was

closed with #1 Vicryl (Serag-Wiesner, Naile, Germany). Following fetal immunization, the uterus was returned to the abdominal cavity and the incision was closed as previously described (Mutwiri, et al., *Immunology* (1999) 97:455-461). At day 142 or 143 of gestation all ewes were anaesthetized with Pentobarbital. The fetuses were exposed, and  
5 euthanized prior to tissue collection.

**Serum cortisol assay:** Serum cortisol analysis was performed in the Endocrine Laboratory, Prairie Diagnostic Services, Western College of Veterinary Medicine (WCVN), Saskatoon, Saskatchewan. Serum samples were stored at -70°C until analyzed  
10 using a fluorescence polarization immunoassay (TDx System, Abbott Laboratories, Irving, TX).

**ELISA:** Polystyrene microtiter plates (Immunolon II, Dynatech Laboratories, Gaithersburg, MD) were coated with 0.1 µg of affinity purified tgD per well and  
15 incubated with serially diluted fetal or ewe sera. The mouse anti-bovine IgG1 monoclonal antibody BIG715A (VMRD Inc, Pullman, WA; 1:4000 dilution), alkaline-phosphatase (AP)-conjugated or biotin-conjugated rabbit anti-sheep IgG (Kirkegaard and Perry Laboratories, Gaithersburg, MD; 1:6000 dilution), or AP-conjugated rabbit anti-sheep IgM (Kirkegaard and Perry Laboratories, Gaithersburg, MD; 1:10000 dilution)  
20 were used to detect gD captured sheep Ig. The reaction was amplified using the AP-streptavidin complex (GibcoBRL, Burlington, ON; 1: 2000 dilution) and visualized with p-nitrophenyl phosphate (PNPP, Sigma-Aldrich, Oakville, ON).

**Virus Neutralization test:** After addition of 5% rabbit serum as a source of  
25 complement, twofold serial dilutions of the non heat-inactivated test sera in medium were mixed with ca. 50 TCID<sub>50</sub> BHV-1 strain 108 in a final volume of 100 µl and incubated for 17 hours at 37°C. Thereafter, 100µl of MDBK cells, corresponding to 1-2 x 10<sup>4</sup> cells per well, were added to each sample and cells were incubated for 4 days at 37°C and 5% CO<sub>2</sub>. Cells were assessed for BHV-1 specific cytopathic effect and serum neutralization  
30 titer was indicated as the highest serum dilution to achieve complete virus neutralization.

**Lymphoproliferative response (LPR):** Blood was collected in EDTA-treated vacutainers (Becton Dickonson, Franklin Lakes, NJ) and mononuclear cells were isolated as described previously (Dudler, et al., in *Immunological methods manual*. 1997 (Lefkovits, I., ed.) Vol. 4, pp. 2075-2078 (Academic press, San Diego/London).

- 5 Lymphocytes were isolated from the lymph nodes and used for LPR and antibody secreting cells assay as previously described (Mutwiri, et al., *Immunology* (1999) 97:455-461). Briefly, cell suspensions were cultured in serum-free medium (AIM-V, GibcoBRL) supplemented with 2% fetal bovine serum (GibcoBRL) and  $2 \times 10^{-5}$   $\mu$ M 2-mercaptoethanol (Sigma-Aldrich). LPR were conducted with  $3 \times 10^5$  cells/well
- 10 (microtiter plates, NUNC, Nalgen Nunc International Corporation, Naperville, IL) in a final volume of 200 $\mu$ l medium. Triplicate cultures were stimulated with 100 $\mu$ l purified gD (0.2 $\mu$ g/ml or 2 $\mu$ g/ml), ConA (Sigma-Aldrich; 1 $\mu$ g/ml) or medium alone. After 3 days incubation, the cells were incubated with [methyl- $^3$ H]thymidine (Amersham Pharmacia Biotech Inc, Baie de Urfe, QC) at a concentration of 0.4  $\mu$ Ci/well for 16 hours. Cells
- 15 were harvested following standard liquid scintillation protocols. Proliferative responses were calculated as the mean of triplicate cultures and expressed as a stimulation index (SI = counts per min in the presence of antigen /counts per min in the absence of antigen).

- Antibody secreting cell assay:** The gD-specific antibody-secreting cells were
- 20 detected using a modified ELISPOT assay, as previously described (Mutwiri, et al., *Immunology* (1999) 97:455-461). Briefly, microtiter nitrocellulose filtration plates were coated overnight with 1 $\mu$ g/ml purified tgD. Unbound protein was removed and  $1 \times 10^6$  cells from individual lymph nodes were added to triplicate wells, in a final volume of 200  $\mu$ l culture medium. After 18 hours incubation at 37°C in a humidified atmosphere with
- 25 5% CO<sub>2</sub> the cells were removed and plates were incubated with biotinylated rabbit anti-sheep IgG (H and L chain-specific, Kirkegaard and Perry Laboratories, Gaithersburg, MD; 1:6000 dilution). Subsequently, the plates were incubated with AP-conjugated streptavidin (Jackson Immunoresearch, Lab. Inc, Westgrove, PA; 1:1000 dilution) and then developed with 5-bromo-4-chloro-3-indolyl phosphate/nitroblue tetrazolium
- 30 (BCIP/NBT) insoluble alkaline phosphatase substrate (Sigma-Aldrich). The frequency of gD-specific ASC per  $1 \times 10^6$  cells was calculated by subtracting the number of ASC



detected in wells not coated with antigen from the number of ASC detected in tgD-coated wells. An inverted light microscope was used to count three replicates for each tissue and data presented are mean values for individual lymph nodes.

5

### Example 1

#### Safety of Fetal Nucleic Acid Immunization

In order to test the safety of oral fetal DNA immunization, vaccine delivery into the amniotic fluid in the oral cavity was performed with nine pregnant ewes at day 123 or 124 of pregnancy (148 day gestation). Four fetuses were immunized with a single  
10 injection of 500 µg tgD-encoding plasmid and four fetuses received PBS. One fetus was immunized with  $5 \times 10^7$  plaque forming units (p.f.u.) of inactivated bovine herpesvirus-1 (BHV-1). Half of the BHV-1 was injected subcutaneously and half was injected into the amniotic fluid in the oral cavity. Following surgery, none of the ewes displayed clinical signs of illness and there were no fetal deaths or abortions. Fetal lambs were of normal  
15 size and displayed normal organ development when tissues were collected at day 145 or 146 of gestation. Histological examination did not reveal pathological changes in any of the fetuses.

To further investigate the effect of DNA vaccination on the fetal development, we measured the level of cortisol in fetal serum. During normal gestation, the level of fetal  
20 serum cortisol begins to increase after day 125 of gestation and a high cortisolemia, shortly before birth, signals the initiation of parturition. In our experiment, all nine fetuses had a high cortisolemia (range = 62 nmol/l to 461 nmol/l) when sera were collected. The normal serum cortisol levels in adult ewes range from 15 to 30nmol/l (Rawlings, et al., *J Toxicol Environ. Health* (1998) 54:21-36). These observations  
25 support the conclusion that *in utero*-DNA vaccination had no adverse effects on fetal development.

Example 2Induction of an Immune Response by Fetal Nucleic Acid Immunization

In order to determine whether *in utero*, oral nucleic acid immunization was capable of producing an immune response, the following experiment was done.

- 5 In particular, to investigate the induction of a gD-specific antibody response by tgD-plasmid immunization, all sera were analyzed with a gD-specific ELISA, described above in the methods. All DNA-vaccinated fetuses (4/4) responded to the single *in utero*-immunization with high titers of gD-specific serum antibodies (Table 1). In contrast, none of the PBS-injected fetuses had detectable levels of gD-specific serum antibodies.
- 10 The fetus immunized with inactivated BHV-1 developed moderate gD-specific titers (Table 1). These observations confirmed that fetal lambs responded to a single DNA vaccination administered via the amniotic fluid.

- The syndesmochorial placenta of sheep excludes maternal immunoglobulin from the fetal circulation (Hein, W.R., *The Immunologist* (1995) 3:12-18). However, to
- 15 substantiate that the gD-specific antibodies were of fetal origin, sera from all ewes were also analyzed in the same ELISA. Maternal gD-specific antibodies were below the level of detection in all ewes whose fetuses were immunized with gD-plasmid. Thus, there was no evidence of antibody transmission from mother to fetus. These results also clearly demonstrate that there was no antibody transfer from the fetus to the mother.

20

Example 3Isotype-switching and Neutralizing Activity

- The hallmark of a functional immune system with cognate B cell-T cell interactions, is isotype-switching from IgM to IgG (Van-den-Eertwegh, et al., *J. Exp.*
- 25 *Med.* (1993) 178:1555-1565; Allen, et al., *Science* (1993) 259:990-993). Monoclonal antibodies specific for bovine IgG1 were used to identify the isotype of the fetal gD-specific antibodies. High titers of gD-specific IgG1 were present only in sera from DNA-vaccinated fetuses (Table 1). To further evaluate the functional significance of the fetal antibody response, all sera were analyzed for virus neutralizing activity. Sera from all
- 30 DNA-vaccinated fetuses contained moderate levels of BHV-1-neutralizing antibodies (Table 1). Furthermore, the level of neutralizing antibodies present in the sera of the

DNA-vaccinated fetuses exceeded the virus neutralizing activity of the serum from the fetus immunized with inactivated BHV-1 (Table 1). As expected from the studies described above, viral neutralizing antibodies were not detected in sera from the PBS-injected fetuses or from the ewes carrying fetuses vaccinated with plasmid DNA.

5

#### Example 4

##### Induction of Cellular Immune Responses in the Blood

To analyze if nucleic acid immunization or fetuses orally *in utero* induced a cellular immune response, the following experiment was conducted. In particular, to determine if immunization with the tgD-encoding plasmid also induced a cell-mediated immune response, mononuclear cells were isolated on day 145 or 146 of gestation from fetal blood to assay gD-specific lymphoproliferative responses (LPR). Lymphocytes from all DNA-vaccinated fetuses proliferated in response to two different concentrations of recombinant gD-protein (0.1 µg/ml and 1.0 µg/ml; Figure 1A). The average stimulation indices (SI) for the DNA immunized group were 5.2 (0.1 µg gD) and 9.7 (1.0 µg gD) respectively, whereas the PBS group had average SIs of 1.1 (0.1 µg gD) and 3.2 (1.0 µg gD). The inactivated BHV-1 immunized fetus had SIs of 1.0 (0.1 µg gD) and 1.7 (1.0 µg gD) respectively. Although, there was not a statistically significant difference between the LPR of DNA- and control-groups, this could be explained by the high LPR of DNA-vaccinated Fetus #7, which had stimulation indices (SI) of 9.8 (0.1 µg gD) and 24.2 (1.0 µg gD) respectively. High levels of serum cortisol can depress *in vitro*-lymphoproliferative responses (Auphan, et al., *Science* (1995) 270:286-290). Thus, the observed magnitude of the LPR of the DNA-immunized fetuses is even more remarkable and may present a minimal estimate of specific immune activation.

25

#### Example 5

##### Induction of a Local Immune Response by Nucleic Acid

##### Immunization into the oral cavity

An important factor for preventing neonatal diseases is the induction of effective mucosal immunity. In order to test for mucosal immunity, a variety of immune assays were used to analyze gD-specific immune responses of lymphocytes isolated from lymph

30



nodes draining various mucosal sites. Two DNA-vaccinated fetuses had numerous gD-specific antibody secreting cells (ASC; ca. 40 gD-specific cells/ million isolated cells) in the retropharyngeal lymph nodes, which drain the oral and nasal cavity (Figure 2). In contrast, no gD-specific ASCs were detected in the retropharyngeal lymph nodes of the BHV-1 immunized and PBS-treated fetuses. No gD-specific ASC were detected in either the mediastinal (lung) or the mesenteric lymph nodes (intestine) of any of the fetuses (data not shown). As expected, low numbers of ASC were identified only in the right, but not the left, prescapular lymph node of the BHV-1 immunized fetus. The right prescapular lymph node drained the site of the subcutaneous BHV-1 injection (Figure 2).

The LPR assay confirmed, that the retropharyngeal lymph node may represent the primary site of immune induction in DNA vaccinated fetuses (Figure 1B). There was a statistically significant difference between the LPR of DNA-vaccinated fetuses when compared with the LPR of PBS-treated fetuses ( $p = 0.015$ ). No gD-specific proliferative responses were detected with cells isolated from the lymph nodes draining other mucosal sites. Taken together, the results from ASC- and LPR-assays clearly demonstrate that injecting gD-plasmid into the amniotic fluid in the fetal oral cavity induced a local immune response in the lymph node draining the site of injection.

#### Example 6

#### Immune Response Following Boost at Birth

In order to determine whether the immune response achieved using *in utero* fetal immunization could be enhanced by a booster administration, the following study was conducted. Four groups of animals were used in this study. Groups I and II were immunized *in utero* with 500  $\mu$ g tgD-encoding plasmid, as described above. Groups III and IV were immunized *in utero* with PBS (no plasmid DNA) as described above. Groups II and IV were subsequently boosted at birth with 500  $\mu$ g tgD-encoding plasmid i.d. Blood was collected at 2 week intervals and serum IgG ELISA titers and PBMC were examined as measures of humoral and cellular immunity, respectively.

As shown in Figures 3 and 4, booster immunization at birth significantly increased the magnitude of the induced immune response.



In conclusion, the present results demonstrate, for the first time, that a single *in utero* immunization with plasmid-DNA effectively induces both mucosal and systemic immune responses in fetal lambs. Moreover, the immune response can be boosted at birth to increase the magnitude of the initial response. Thus, fetal DNA vaccination, with  
5 or without a subsequent boost, provides a safe and effective method for preventing or reducing the high risk of vertical disease transmission during pregnancy and following birth. Vaccination may be performed using surgical techniques such as amniocentesis, which involves minimal surgical invasion. The present observations have significant application for developing new gene therapy strategies to target the upper and lower  
10 respiratory tract. The present techniques may also be used for targeting gene expression to various mucosal sites where transient expression of a plasmid DNA may be of significant value.

9000-0058 PATENT

5

Table 1: gD-specific antibody response in fetal sera

Fetus	Immunogen	Total gD-specific antibody titre*	gD-specific IgG1 antibody titre <sup>+</sup>	Neutralizing antibody titre
1	PBS	11 ± 15	nd <sup>++</sup>	< 2
2	PBS	14 ± 20	nd	< 2
3	PBS	10 ± 14	nd	< 2
4	PBS	11 ± 16	nd	< 2
5	tgD-plasmid	2047 ± 253	1586 ± 120	32
6	tgD-plasmid	5034 ± 333	3864 ± 233	64
7	tgD-plasmid	2628 ± 258	4948 ± 799	32
8	tgD-plasmid	1146 ± 672	412 ± 267	8
9	irr. BHV-1	666 ± 125	1202 ± 334	< 2

\* Rabbit anti-sheep IgG (H+L chain-specific)

+ B1g715A monoclonal antibody

++ not detected

Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720